



Liquid Miscellaneous Secondary Nutrients and
Micronutrients Fertilizer (Serial Number 4-42)
Fertilizer Registration Certificate Number:0727085

Root FORTE

速益根 110

SPECIAL FOR ROOTING AND GERMINATION

- Promote root growth
- Strengthen the vigor of new sprouts
- Activation of mesophyll tissue
- Delaying crops aging
- Enhance the soil aggregate structure

Active ingredients

Total nitrogen	1.5 %
Water-soluble potassium oxide	1.5 %
Water-soluble calcium oxide	3.0 %
Water-soluble boron	0.25 %
Water-soluble molybdenum	0.010 %

Method of Use

All Crops

1. Seedling stage: After 2 new leaves have grown, dilute 1000 times for spraying leaves or root watering. The frequency of use depends on the growth of the plant.
2. After transplanting (about 5~7 days): dilute 500 times for root watering. The frequency of use depends on the crop's growth status.

**Do not use this product in sunlight and high temperature environment; avoid transpiration effects to affect plant growth.

**When spraying the leaves, be sure to dilute the solution according to the specified dilution ratio to prevent fertilizer burns on leaves.

**The used amount for spraying or watering after dilution can be adjusted according to the weather, soil moisture, and plant growth status.

Cautions

*This product contains Boron (Br) and Molybdenum (Mo), and excessive use may cause toxicity and soil pollution. Do not spray on edible parts one week before harvesting.

*Please apply fertilizer according to the recommended method and usage amount.

*Shake well before use.

*Keep in a cool and dry place away from direct sunlight.

*When this product is mixed with other materials, perform a small-scale test first.



Made in Taiwan

Manufacturer:

Anji Biotech Corp.

Distributor: KWAN STAR CO., LTD.

21F-1, No.268, SEC. 1, WEN HWA RD., BANCIAO DIST.,
NEW TAIPEI CITY, TAIWAN 22041

TEL:886-2-89690978

business@kwanstar.com



LINE
@kwanstar



FB



WEB



Shopee

CROP RECOMMENDATIONS FERTILIZER PROGRAM

DOSAGE, LITER/HECTARE

1LTR/2HA
(FOLIAR APPLICATION)

1. Diluted 1000 times.
2. For Foliar application or root watering.

VEGETABLES, IN GENERAL (SPRING ONION, CELERY, LEEK, BASIL, CORIANDER)



1LTR/2HA

At 2-3 leaves - after transplanting
(the frequency of use depends on the growth status)

FRUIT-BEARING FLOWERING CROPS (EGGPLANT, BELL PEPPER, CHILI PEPPER, TOMATO, WOLFBERY)



1LTR/2HA

At 2-3 leaves, 1 time
(FOLIAR APPLICATION)

VEGETABLES, IN GENERAL (CABBAGE, LETTUCE)



1LTR/2HA

At 2-3 leaves
(the frequency of use depends on the growth status)

CAULIFLOWER, BROCCOLI



1LTR/2HA

At 2-3 leaves - after transplanting
(the frequency of use depends on the growth status)

BEANS, IN GENERAL (PEAS, EDAMAME, RED BEANS, GREEN BEANS, SOYBEANS, BLACK BEANS)



1LTR/2HA

At 2-3 leaves
(the frequency of use depends on the growth status)

PEANUTS



1LTR/2HA

At 2-3 leaves
(the frequency of use depends on the growth status)

CORN



1LTR/2HA

At 2-3 leaves, 1 time
(FOLIAR APPLICATION)

CUCURBITS, MELONS



1LTR/2HA

At 5 leaves, 1 time
(FOLIAR APPLICATION)

SOFT FRUITS (STRAWBERRY)



1LTR/2HA

At 2-3 leaves, 1 time
(FOLIAR APPLICATION)

TEA TREES



1LTR/1HA

Growing more than three years, after harvest,
waiting for the new leaves have grown
New leaves fully expansion, every 30 days
(FOLIAR APPLICATION)

ORCHARDS



N/A

Anytime, (every 14 days)
(FOLIAR APPLICATION)
Applying can be adjusted according to the size
of the fruit trees

BANANA



N/A

Anytime, (every 14 days)
(FOLIAR APPLICATION)
Applying can be adjusted according to the size of